



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

517 (Geometry). Proposed by R. P. BAKER, University of Iowa.

The coördinates of the vertices of a regular icosahedron can be expressed rationally in terms of $(\sqrt{5} - 1)/4$ and $\sqrt{(5 + \sqrt{5})}/8$, that is, $\cos(2\pi/5)$ and $\sin(2\pi/5)$. Prove (1) that the cosine alone is sufficient; (2) that the irrationalities cannot be reduced further. (The theorem that they cannot be rational is proved in books on crystal theory.)

SOLUTION BY C. F. GUMMER, Queen's University, Kingston.

Since the coördinates of the center are expressible rationally in terms of the vertices, we may suppose it taken as origin. Moreover the vertices are all rational in terms of any three adjacent ones A, B, C ; for a fourth is the reflection of A for the plane OBC , and so on. Suppose first that A, B, C have coördinates (o, o, r) , $(r \sin \alpha, o, r \cos \alpha)$, $(r \sin \alpha \cos(2\pi/5), r \sin \alpha \sin(2\pi/5), r \cos \alpha)$. Since $AB = BC$, we find $\cos \alpha = 1/\sqrt{5}$; and α is the angle subtended at O by an edge. Hence, the general equations for A, B, C (with the origin for center) are

$$\begin{aligned} x_1^2 + y_1^2 + z_1^2 &= x_2^2 + y_2^2 + z_2^2 = x_3^2 + y_3^2 + z_3^2 = \sqrt{5}(x_1x_2 + y_1y_2 + z_1z_2) \\ &= \sqrt{5}(x_2x_3 + y_2y_3 + z_2z_3) = \sqrt{5}(x_3x_1 + y_3y_1 + z_3z_1). \end{aligned}$$

Evidently no rational solution exists, which proves the second part of the theorem.

A simple solution is found by assuming $x_1 = y_2 = z_3 = u$, $x_2 = y_3 = z_1 = v$, $x_3 = y_1 = z_2 = w$, so that the equations become $u^2 + v^2 + w^2 = \sqrt{5}(uv + vw + wu)$. If $w = 0$, $u^2 + v^2 = \sqrt{5}uv$, which is satisfied by $u = 1$, $v = 2 \cos(2\pi/5)$. Therefore, A, B, C may be taken to be $(1, 0, 2 \cos(2\pi/5))$, $(2 \cos(2\pi/5), 1, 0)$, $(0, 2 \cos(2\pi/5), 1)$.

518 (Geometry). Proposed by ROGER A. JOHNSON, Cleveland, Ohio.

If one angle of a triangle is 60° , the Euler line (the line through the circumcenter, orthocenter, and median point) is perpendicular to the bisector of that angle; and if one angle is 120° , the Euler line is parallel to the bisector of that angle.

SOLUTION BY J. L. COOLIDGE, Harvard University.

Let the vertices of the triangle be A_1, A_2, A_3 , the middle points of the sides M_1, M_2, M_3 , the feet of the altitudes H_1, H_2, H_3 , the circumcenter O , and the orthocenter H .

Let us take A_1 as the angle in which we are interested, and assume it first to be acute.

The angle which the external bisector forms with A_2A_3 is $\frac{1}{2}(A_2 - A_3)$ and its tangent

$$\tan \frac{1}{2}(A_2 - A_3) = \frac{1 - \cos(A_2 - A_3)}{\sin(A_2 - A_3)}.$$

The tangent of the angle which the Euler line makes with A_2A_3 is

$$\frac{OM_1 - HH_1}{A_2M_1 - A_2H_1}.$$

We have

$$OM_1 = r \cos A_1 = r[\sin A_2 \sin A_3 - \cos A_2 \cos A_3],$$

$$A_2M_1 = r \sin A_1 = r[\sin A_2 \cos A_3 + \cos A_2 \sin A_3],$$

$$HH_1 = (A_2H_1) \cot A_3 = 2r \cos A_2 \cos A_3,$$

and

$$A_2H_1 = a_3 \cos A_2 = 2r \cos A_2 \sin A_3.$$

The tangent of this angle is

$$\frac{\sin A_2 \sin A_3 - 3 \cos A_2 \cos A_3}{\sin(A_2 - A_3)}.$$

The two tangents are equal if

$$1 - \cos A_2 \cos A_3 - \sin A_2 \sin A_3 = \sin A_2 \sin A_3 - 3 \cos A_2 \cos A_3, \text{ or } \cos(A_2 + A_3) = -\frac{1}{2}.$$

Hence,

$$A_2 + A_3 = 120^\circ.$$